

CONTACT:

Maggie Moon, MS, RD
Pollock Communications
mmoon@pollock-pr.com
(212) 941-1414

TEA AND HEALTH

-- An Overview of Research on the Potential Health Benefits of Tea --

INTRODUCTION

Tea is an ancient beverage steeped in history and romance and loved by many. In fact, so popular is tea that it is the most commonly consumed beverage in the world after water. Although tea had a modest beginning (it was discovered by accident), its popularity spread from its origins in China to Western Europe and the Americas. Throughout history, tea has been believed by many to aid the liver, destroy the typhoid germ, purify the body and preserve mental equilibrium. Over the past few decades, scientists have taken a closer look at the potential health benefits of tea and have discovered that much of the folklore about tea may actually be true.

HOW TEA WORKS IN THE BODY

Tea contains flavonoids, naturally occurring compounds that have been shown to have antioxidant properties. Antioxidants work to neutralize free radicals, which scientists believe, over time, damage elements in the body, such as genetic material and lipids, and contribute to many chronic diseases. Recent research has explored the potential health attributes of tea through studies in humans and animal models, and through *in vitro* laboratory research. For the most part, studies conducted on Green and Black Tea, which are both from the *Camellia sinensis* plant, have yielded similar results.

An examination of recently released dietary intake data from the National Health and Nutrition Examination Survey (NHANES) reported that differences in total flavonoid intake among subgroups are principally associated with the percentage of tea consumers and their prevalence of tea consumption.¹ Valuable new information has also been recently reported regarding the bioavailability and metabolism of tea flavanols using novel approaches with *in vitro* digestion models as well as in human studies.^{2,3} These categories of data are important as individuals consider what foods to include to increase bioavailable and bioactive phytochemicals.

Recent research suggests that tea and tea flavonoids may play important roles in various areas of health and may operate through a number of different mechanisms still being explored. Recent findings about tea and health include:

- A recent review of tea and health in the elderly suggested that there is compelling evidence for the efficacy of tea in benefiting cardiovascular disease.⁴ The antioxidant properties of tea flavonoids may play a role in reducing the risk of cardiovascular disease by decreasing lipid oxidation,⁵ reducing the instances of heart attacks and stroke,^{6,7} and may beneficially impact blood vessel function,⁸ an important indicator of cardiovascular health.
- Tea flavonoids may lower the risk of certain cancers by inhibiting the oxidative changes in DNA from free radicals and some carcinogens.⁵ Tea may also promote programmed cell death, or apoptosis,⁹ and inhibit the rate of cell division, thereby decreasing the growth of abnormal cells.⁵
- Tea polyphenols are bioavailable to the brain and can act via antioxidant, iron-chelation, signal transduction modulation, and other mechanisms to effect neuroprotective and/or neurorescue action, with potential implications for age-related dementia, Alzheimer's and Parkinson's diseases.¹⁰
- A unique tea amino acid, L-theanine (γ -glutamylethylamide), plays a role in attentional processing in synergy with caffeine.¹¹
- Tea-drinking has been associated with oral health¹² and bone health.¹³
- Compounds in tea other than flavonoids have been shown to support the human immune system.¹⁴
- Due to the substantial data documenting tea's health benefits, recently published *Healthy Beverage Guidelines* suggest water, tea and coffee should provide the majority of daily fluid intake.

Unsweetened Tea (up to eight servings per day) is recommended because it is virtually calorie-free, delivers antioxidant phytonutrients and has less caffeine than coffee (about 40 mg per serving).¹⁵

TEA'S ROLE IN CARDIOVASCULAR HEALTH

Human population studies have found that people who regularly consume three or more cups of Black Tea per day have a reduced risk of heart disease and stroke. Clinical studies suggest that the risk reduction associated with Black Tea consumption may be due to improvement in some risk factors for cardiovascular disease, including blood vessel function, platelet function and a reduction in oxidative damage.

While researchers are still examining the various mechanisms by which tea flavonoids function, some studies suggest multifunctional mechanisms, meaning that several mechanisms work in tandem to collectively improve markers for cardiovascular health. Important areas of tea and cardiovascular health research include blood vessel and endothelial function, or the ability of the blood vessels to dilate to allow for proper blood flow, serum cholesterol levels and Low Density Lipoprotein (LDL) cholesterol oxidation. Each of these factors impact the risk of myocardial infarctions (heart attacks), stroke and cardiovascular disease. Study findings in the area of tea and the reduction in cardiovascular disease risk include the following:

Cardiac Events

- A total of 3,430 men and women aged 30-70 years from the Saudi Coronary Artery Disease Study were examined and 6.3 percent were found to have indications of coronary heart disease (CHD). The researchers found that those who drank more than six cups of tea per day (>480 mL) had significantly lower prevalence of CHD than non-tea drinkers, even after adjustment for risk factors like age and smoking.¹⁶ The researchers also found that drinking six or more cups of Black Tea per day was associated with decreased serum cholesterol and triglyceride concentrations.
- The Zutphen study, which assessed 805 male subjects over a period of five years, found that the incidence of fatal and nonfatal first myocardial infarction and mortality from stroke decreased significantly as intake of flavonoids, derived mainly from tea, increased in a dose-dependent manner.⁶ A follow-up to this study found that high intake of flavonoids significantly lowered the risk of stroke in study participants.⁷
- A Harvard study examined 340 men and women who had suffered heart attacks and compared them to matched control subjects. They found that those who drank a cup or more of Black Tea daily had a 44 percent reduction in the risk of heart attack compared to non-tea drinkers.¹⁷
- Another recent Harvard study of 1,900 people found that those who consumed tea during the year prior to a heart attack were up to 44 percent more likely to survive over the three to four years following the event. Those who consumed fewer than 14 cups of tea per week experienced a 28 percent reduced death rate, and those who consumed more than 14 cups of tea per week were found to have a 44 percent reduced death rate, as compared to non-tea drinkers.¹⁸
- Dutch researchers assessed 4,807 subjects aged 55 years or older without prior history of heart attack. After a four to seven year follow up period, the researchers determined that those who drank three or more cups of tea per day (375mL) were 43 percent less likely to develop myocardial infarction and 70 percent less likely to die from myocardial infarction than non-tea drinkers.¹⁹
- A recent meta-analysis discovered that consumption of three cups of tea per day was associated with an estimated decrease of 11 percent in the incidence of myocardial infarction (heart attack).²⁰
- A large Japanese population study of over 40,000 middle-aged Japanese reported that, among men and women, those who drank just over two cups (about 17 ounces) of Green Tea per day reduced their risk of death from cardiovascular disease by 22 to 33 percent, compared to those who drank less than a half-cup (3.5 ounces) of Green Tea daily.²¹

Cholesterol Reduction

- Researchers from the United States Department of Agriculture (USDA) studied the effect of tea on 15 mildly hypercholesterolemic adult participants following a "Step 1" type diet moderately low in fat and cholesterol, as described by the American Heart Association and the National Cholesterol Education Program. After three weeks, researchers found that five servings of Black Tea per day reduced LDL ("bad") cholesterol by 11.1 percent and total cholesterol (TC) by 6.5 percent compared to placebo beverages.²² Recent clinical trials have not confirmed these results, however additional work is being done in this area.
- **The mechanism behind the blood cholesterol lowering effects of tea may be rooted in the effect of theaflavins, through interfering with the formation of dietary mixed micelles, which could result in**

reduced intestinal cholesterol absorption. Theaflavin-treated micelles/particles were analyzed and theaflavins were shown to have a dose-dependent inhibitory effect on the incorporation of cholesterol into micelles. The primary theaflavin identified for its effects was theaflavin-3-gallate.²³

Tea and Cardiovascular Disease (CVD) Risk Factors

- Dutch researchers found that study participants who drank one to two cups of Black Tea daily had a 46 percent lower risk of severe aortic atherosclerosis, a strong indicator of cardiovascular disease. Those who drank more than four cups of tea a day had a 69 percent lower risk.²⁴
- A recent clinical study showed that short- and long-term consumption of Black Tea by subjects with coronary artery disease restored endothelial and blood vessel function to levels similar to that of healthy subjects.²⁵ Endothelial function is the ability of the inner lining of blood vessels to dilate in response to increased blood flow.
- Another clinical study found that regular ingestion of tea resulted in a significant and consistent increase in endothelium-dependent and endothelium-independent blood vessel dilation.²⁶ Subjects with mild elevations in serum cholesterol or triglyceride concentrations consumed either five cups of Black Tea per day for four weeks or hot water. The researchers hypothesized that one mechanism for the apparent beneficial effects of tea on cardiovascular health could be this improved vasodilator function.
- An *in vitro* study found that Green Tea polyphenols inhibit the proliferation of aortic smooth muscle cells to prevent the development of atherosclerosis.²⁷
- *In vitro* studies have shown that tea flavonoids protect low-density lipoproteins from oxidation, inhibit plasma lipid peroxidation, platelet aggregation and thromboxane formation – all factors important for maintaining a healthy circulatory system.^{28,29} Studies in animals are promising, but human studies conducted to date on the effect of tea consumption on LDL oxidation are inconclusive.
- A double-blind crossover design clinical study found that Black Tea versus a caffeinated control beverage improved coronary flow velocity reserve, a measure of increased blood flow in vessels.³⁰
- Animal model studies found that Green or Black Tea both improved risk factors for CVD, including reduced blood lipids, serum antioxidants levels and improvements in blood clotting factors.³¹
- A longitudinal study with more than 1,500 Taiwanese found that those who drank more than 2 1/2 cups of tea per day reduce their risk of developing high blood pressure by 65 percent compared to those who drank the least amount of tea.³²

TEA'S ROLE IN CANCER RISK REDUCTION

Preliminary research suggests that the flavonoids in tea could play a role in human cancer risk reduction possibly by combating free radical damage, inhibiting uncontrolled cell growth (cell proliferation), by promoting programmed cell death (apoptosis) and boosting the immune system to help fend off the development and promotion of cancer cells. Leading scientists worldwide are actively studying these potential mechanisms, and clinical trials and population studies are underway. More evidence is needed before any definitive conclusions can be drawn. Recent findings include:

DNA Damage

- Oxidative DNA damage is implicated in the development of various forms of cancer. Recent studies have found that smokers who drank four cups of decaffeinated Green Tea per day demonstrated a 31 percent decrease in biomarkers of oxidative DNA damage in white blood cells as compared to those who drank four cups of water after four months (P=0,001).^{33,34}
- Epigallocatechin gallate (EGCG) may protect normal cells from cancer-causing hazards as well as eliminate cancer cells through apoptosis. Researchers tested the potential anti-cancer benefits of Green Tea polyphenol, EGCG, in hamster cells and discovered that EGCG suppressed DNA changes and damage from carcinogens. EGCG also protected from further damage from the carcinogens and inhibited growth and multiplication of cancer cells.³⁵
- An animal study identified beneficial changes in immune function after Black Tea ingestion in cancer-bearing animals. Black Tea beneficially altered immune responses that helped protect immune cells against harmful cancerous cells. The study found Black Tea acted like anti-cancer drugs that help boost the immune system without promoting the proliferation of cancerous cells.³⁶

Digestive Cancers

- An epidemiological study conducted by the University of North Carolina found consumption of the equivalent of 2.5 cups of tea per day or more was associated with a 60 percent drop in rectal cancer risk among Russian women from Moscow, as compared to women who drank relatively less than 1.2

cups of tea per day. Those women who drank approximately 1.2 to 2.5 cups of tea per day had a 52 percent reduction in the risk of rectal cancer.³⁷

- Based on data from the NHANES I Follow-Up study (NHEFS), researchers found that tea drinkers had about a 42 percent reduced risk of colon cancer as compared to non-tea drinkers. Men who drank more than 1.5 cups of tea per day were found to have a 70 percent lower colon cancer risk.³⁸
- Researchers who followed a group of over 34,000 postmenopausal healthy women between 55 – 69 years of age for 12 years found that those consuming high levels of catechins experienced up to a 45 percent decrease in the instances of rectal cancer. Catechins are a class of flavonoids found in tea, fruits and vegetables. Catechins derived from tea were most strongly linked to a decrease in rectal cancer.³⁹
- The Iowa Women's Study, which followed post-menopausal women between the ages of 55 and 69 for eight years, found that participants who drank two or more cups of tea per day had a 32 and 60 percent reduced risk of developing digestive and urinary tract cancers, respectively.⁴⁰
- A study conducted with members of the Shanghai Cohort (18,244 men aged 45-64 years at recruitment with up to 12 years of follow-up) discovered a statistically significant inverse relationship between positive tea polyphenol levels (as measured in urine) and gastric cancer.⁴¹
- A large population-based case-control study found an inverse relationship between Green Tea consumption and the risk of colon, rectal and pancreatic cancer. Male participants, who drank the equivalent of 4.5 servings of tea per day, had an 18 percent decrease in colon cancer risk and 28 percent decreased risk of rectal cancer. Female participants, who drank 3 servings of tea per day, were observed to have a decreased risk of colon and rectal cancer by 33 percent and 43 percent, respectively. Risk of pancreatic cancer was also reduced in both men and women by 37 percent and 47 percent, respectively.⁴²
- Researchers examined whether a combination of two compounds known to exhibit anti-cancer activity, Green Tea polyphenol, EGCG, and sulindac (a non-steroidal anti-inflammatory drug), would work synergistically to prevent colon cancer carcinogenesis in rats. Findings suggested that EGCG and sulindac worked together to suppress pre-cancerous lesion formation by enhancing programmed cell death (apoptosis).⁴³
- Researchers sought to investigate the effect of Black Tea polyphenols (BTP) on induced DNA damage to colon mucosa in an animal model. Findings suggest that induced DNA damage to the colon mucosa is prevented by consumption of Black Tea polyphenols.⁴⁴
- Major compounds of Green and Black Tea, EGCG and theaflavins respectively, are known to inhibit proteins which are closely associated with tumor growth and metastasis. These polyphenols exhibited apoptosis-inducing activity for human colon cancer cell lines.⁴⁵
- Researchers in Taiwan discovered a link between EGCG and cancer risk reduction. The researchers found that the Green Tea polyphenol inhibited proliferation of the cancer cells by inducing cell death and blocking cell cycle progression.⁴⁶
- Researchers at the University of South Carolina used animals with colon cancer and provided them with either water or Green Tea as their beverage. They subsequently found that those given Green Tea significantly reduced their risk of developing new colon cancer tumors, suggesting that Green Tea was effective in the initial stages of colon carcinogenesis.⁴⁷

Prostate Cancer

- Researchers at the University of Wisconsin, Madison reviewed the existing literature about tea as a preventative measure for prostate cancer among men. Based on epidemiological, in vitro and in vivo studies, the researchers suggest that tea—especially Green Tea—may be a good public health recommendation that may help prevent prostate cancer.⁴⁸

Skin Cancer

- According to a study conducted by the University of Arizona, participants who drank iced Black Tea and citrus peel had a 42 percent reduced risk of skin cancer.⁴⁹
- Hot Black Tea consumption is associated with a significantly lower risk of squamous cell carcinoma (SCC), a form of skin cancer; tea concentration (strength), brewing time and temperature all influence the potential protective effects of hot Black Tea on SCC.⁵⁰
- Oral consumption of Green or Black Tea decreased the number of tumors in mice following exposure to UV radiation.⁵¹
- Green Tea polyphenols may have cancer prevention potential, especially in the case of solar UV-induced cancer.⁵²
- Research suggests that compounds in Green Tea may protect skin from ultraviolet (UV) radiation-induced damage when applied topically.⁵³

- Topical treatment of Green Tea polyphenols on human skin prior to UV exposure inhibited indicators of DNA damage, thus inhibiting photocarcinogenesis, or UV-induced skin cancer.⁵⁴
- Experiments that show that administration of Green Tea, Black Tea or specific flavonoids in tea inhibited the growth of established nonmalignant and malignant skin tumors in tumor-bearing mice. In addition, oral administration of Black Tea inhibited DNA synthesis and enhanced cell death (apoptosis) in both nonmalignant and malignant tumors in tumor-bearing mice.⁵⁵

Oral Cancers

- A human intervention trial examined the effect of treating superficial precancerous lesions (leukoplakia) in the mucosal lining of the mouth with a mixed tea product. After the six-month trial, partial regression of the lesions was observed in 37.9 percent of the group treated with tea as compared to only 10 percent of those treated with a placebo.⁵⁶
- Researchers examined the effects of tea and curcumin, a spice and food-coloring agent, on oral cancer in hamsters. Hamsters were treated with a topical cancer-causing solution inside the cheek three times a week for six weeks. Two days after the last treatment of the solution, the hamsters were given Green Tea as drinking fluid or curcumin applied topically three times per week, the combination of Green Tea and curcumin treatment, or no treatment for 18 weeks. At the end of this period, the scientists observed that the combination of tea and curcumin significantly decreased the number of visible tumors and tumor volume. Furthermore, tea alone and in combination with curcumin increased cancer cell death (apoptosis).⁵⁷

Lung Cancer

- In a recent review of observational studies on tea consumption, flavonoid intake, and lung cancer risk, evidence suggests beneficial associations for green and black tea, especially among never-smokers. The review notes that studies reporting increased risk with high tea intake are older and later published data have not confirmed these concerns.⁵⁸
- Studies comparing groups of mice treated with a tobacco-specific carcinogen and receiving either water or water enriched with tea-derived antioxidants found that the tea-fed mice developed 24 percent fewer lung tumors and the average size of the tumors was 38 percent smaller as compared to the water-fed mice.^{59,60}
- Tea catechins were evaluated for their effects on cell proliferation, apoptosis and associated gene expression in highly metastatic human lung cancer cells. A significant reduction in cell proliferation after exposure to tea catechins was noted. It is suggested that tea compounds can influence genetic alteration to reduce the growth and survival of human lung cancer cells.⁶¹

Ovarian Cancer

- A case-control study conducted in China, which employed 254 patients with histologically confirmed epithelial ovarian cancer and 652 control subjects, determined tea consumption based on a validated questionnaire and found that, after accounting for demographic, lifestyle and familial factors, ovarian cancer risk declined with increasing frequency and duration of overall tea consumption.⁶²
- A population-based study involving over 61,000 Swedish women aged 40-76 found that drinking Black Tea was associated with a reduced risk of ovarian cancer. The study found that women who drank the most tea—green or black—were least likely to develop ovarian cancer over the 15-year study follow-up. Women who drank two or more cups of tea daily experienced a 46 percent reduction in risk compared to women who reported not drinking tea. Even small amounts of tea (less than one cup per day) reduced risk by 18 percent, while one cup per day reduced risk by 24 percent. Although previous studies evaluating the effects of tea consumption and ovarian cancer found inconsistent results, the researchers noted that the large size of this study and long-term follow-up provides compelling evidence that tea drinking may indeed offer protection against this type of cancer.⁶³

Breast Cancer

- Results from a large epidemiologic study examining the association of regular tea consumption with the risk of breast cancer found that women under 50 years with moderate tea consumption (three or more cups per day) had a 37% reduced breast cancer risk.⁶⁴

TEA AND NEUROLOGICAL DECLINE

- A recent human study examined the effect of the unique tea amino acid L-theanine (γ -glutamylethylamide) on attention-related task performance. Task performance was measured by electroencephalographic (EEG), or the measurement of [electrical](#) activity produced by the [brain](#) as

recorded from [electrodes](#) placed on the scalp. The results suggest L-theanine plays a role in attentional processing in synergy with caffeine.⁶⁵

- In an animal model study, Japanese researchers found that theanine, the amino acid present virtually only in tea, may help prevent memory declines as we age by decreasing neuronal cell death. In their study, animals who were given theanine and were then subjected to repeated memory impairment, had less memory damage to their brains compared to animals who did not receive theanine.⁶⁶
- Newly published research reports that tea polyphenols, particularly (–)-epigallocatechin-3-gallate, are bioavailable to the brain and can act via antioxidant, iron-chelation, signal transduction modulation, and other mechanisms to effect neuroprotective and/or neurorescue action, with potential implications for age-related dementia, Alzheimer's and Parkinson's diseases.⁶⁷
- In an animal model study, researchers found that the Green Tea catechin EGCG was capable of reducing biomarkers associated with Alzheimer's disease. Using strains of mice transgenically bred to be prone to develop Alzheimer's disease, the researchers found that exposure to EGCG resulted in reduced production of amyloid protein, a marker for the development of plaques associated with Alzheimer's disease.⁶⁸
- A prospective cohort study of nearly 30,000 Finnish adults aged 25 to 74 years old, who were followed for 13 years, found that tea drinking was associated with a reduced risk of Parkinson's disease. Among tea drinkers, those who reported drinking three or more cups of tea per day were 69% less likely to develop Parkinson's disease compared to those who reported not drinking tea.⁶⁹

TEA AND METABOLISM, OBESITY AND BODY COMPOSITION

Preliminary research suggests that drinking tea may have effects on body weight, fat accumulation and insulin activity. While it may be premature to draw firm conclusions based on early research, key findings include the following:

- Green Tea extract was found to significantly increase 24-hour energy expenditure and fat oxidation in healthy men.⁷⁰
- After three months of consumption of Green Tea extract by moderately obese patients, body weight decreased by 4.6 percent and waist circumference decreased by 4.48 percent.⁷¹
- Average fat oxidation rates were 17% higher after acute green tea extract intake (890 +/- 13 mg polyphenols; 366 +/- 5 mg EGCG) during moderate-exercise (30 minutes of cycling at 60% VO₂max) in 12 healthy men, compared to a placebo (p<0.05).⁷²
- Japanese researchers found that in a 12-week, double-blind and placebo-controlled study, green tea catechins led to a reduction in body fat, blood pressure and LDL cholesterol compared to the control group. The authors suggest that Green Tea catechins may help prevent obesity and reduce risk for cardiovascular disease.⁷³
- A follow-up study of the effects of tea catechins on body fat reduction in humans was conducted by examining the effect of drinking Oolong Tea with added Green Tea extract in healthy, moderately overweight men. A double-blind study was performed in which the test subjects ingested one bottle of Oolong Tea containing 690 mg of catechins and control subjects ingested one bottle of Oolong Tea containing 22 mg of catechins for 12 weeks. Researchers found that body weight, body mass index (BMI), waist circumference, body fat mass and subcutaneous fat were all significantly lower in the high catechin-ingesting group as compared to the control group. Measures of LDL cholesterol oxidation were positively associated with the beneficial changes in body fat mass in the high catechin group. Researchers concluded that daily consumption of 690 mg of catechins, the equivalent of five cups of strong Green Tea, might be useful in preventing and improving obesity.⁷⁴
- Researchers examined mice which were fed either a low-fat diet, high-fat diet or high-fat diet supplemented with 0.1-0.5 percent tea catechins for 11 months. The scientists then measured body weight, fat tissue mass and liver fat content and discovered that supplementation with tea catechins resulted in a significant reduction of high-fat diet-induced body weight gain and visceral and liver fat accumulation.⁷⁵
- Researchers compared the body weight and fat mass of mice that were fed a low-fat or high-fat diet, with swimming or not, and with or without tea catechins. They found that, when fed a high-fat diet, tea catechins helped reduce fat accumulation by 18 percent and exercise alone reduced accumulation by 14 percent. However, mice that exercised and had catechins reduced fat accumulation by 33 percent. This evidence suggests that tea catechins may increase fat metabolism, enabling the body to burn more fat as fuel and store less in the body.⁷⁶
- Animals fed a high-fat, high-calorie diet to promote excessive weight gain and obesity were given Green Tea extract or placebo and their energy expenditure and fat oxidation were measured. The researchers found that Green Tea extract alone, as well as when combined with exercise, increased energy expenditure and stimulated fat catabolism. The researchers concluded that Green Tea extract

combined with regular exercise stimulates fat metabolism and may attenuate obesity caused by a high-fat diet more effectively than Green Tea extract or exercise alone.⁷⁷

- Animals fed a diet high in catechin-rich green tea extract were found to increase running times to exhaustion by up to 30 percent compared to a control animal. In addition, Green Tea extract appeared to shift metabolism so that the animals burned body fat and spared muscle glycogen, thereby increasing endurance time to exhaustion.⁷⁸

Diabetes

- Recent USDA research reviewed the effect of tea on insulin sensitivity and risk factors for diabetes. Epidemiologic studies suggest some relation between tea consumption and a reduced risk of type 2 diabetes. Some human clinical studies show tea and its components improved control of glucose metabolism and endothelial function, the decline of which is a marker for vascular diseases.⁷⁹
- Researchers at the United States Department of Agriculture (USDA) conducted a study to examine the insulin-enhancing properties of tea and its components. An *in vitro* test using a fat cell assay found that tea, as normally consumed, increased insulin activity greater than 15-fold. Green, Black and Oolong Tea all yielded insulin-increasing results. The researchers separated the components of the tea using a high-performance liquid chromatography and discovered that several known compounds found in tea were shown to enhance insulin, helping cells recognize and respond to the hormone. The greatest activity was elicited by EGCG followed by epicatechin gallate, tannins, and theaflavins.⁸⁰

TEA'S ROLE IN IMMUNE FUNCTION

Researchers from Brigham and Women's Hospital and Harvard University recently published novel new data indicating that tea contains a component that can help the body ward off infection and disease and that drinking tea may strengthen the immune system.

The researchers identified a substance in tea, L-theanine, which primes the immune system in fighting infection, bacteria, viruses and fungi. A subsequent human clinical trial showed that certain immune cells of participants who drank five cups of Black Tea a day for two to four weeks secreted up to four times more interferon, an important part of the body's immune defense, than at baseline. Consumption of the same amount of coffee for the same duration had no effect on interferon levels. According to the authors, this study suggests that drinking Black Tea provides the body's immune system with natural resistance to microbial infection.⁸¹

TEA'S ROLE IN ORAL HEALTH

- Tea may also contribute to oral health. The flavonoids in tea may inhibit the plaque-forming ability of oral bacteria and the fluoride in tea may support healthy tooth enamel.^{82,83}
- A recent study conducted at the New York University Dental Center examined the effects of Black Tea extract on dental caries formation in hamsters. Compared to those who were fed water with their food, hamsters which were fed water with Black Tea extract developed up to 63.7 percent fewer dental caries.⁸⁴
- Drinking tea is minimally erosive to tooth enamel according to a recent study comparing tea (green and black) to soda and orange juice using in-vitro tests. Water was used as the non-erosive control, and vinegar was the erosive control. The 20-week study was conducted under controlled conditions, and results were categorized as highly, moderately, or minimally erosive. Soda and orange juice were shown to be moderately erosive, and vinegar remained highly erosive.⁸⁵
- Drinking green tea was inversely related to periodontal (gum) disease, per a modest drop in probing depth, clinical attachment loss, and bleeding on probing. The study analyzed 940 Japanese men, aged 49 to 59 years old who took part in a comprehensive health exam. The relationship seems to be dose-dependent. Each additional cup of tea was associated with a greater decrease in gum disease factors.⁸⁶

TEA AND REDUCED RISK OF KIDNEY STONES

Increased intake of fluids is routinely recommended for people who have had kidney stones to reduce the likelihood of recurrence. A recent study that followed 81,093 women for eight years suggests that beverage choice may also affect kidney stones development. The study found that for each eight-ounce cup of tea consumed daily by female participants with no previous history of kidney stones, the risk of developing stones appeared to be lowered by eight percent.⁸⁷ An earlier study of 45,289 men reported a

similar relationship, suggesting that for each eight-ounce serving of tea consumed daily, a 14 percent decrease in risk of stone development was observed.⁸⁸

TEA AND REDUCED RISK OF OSTEOPOROSIS

Although high caffeine intake has been suggested to be a risk factor for reduced bone mineral density (BMD), research indicates that drinking tea does not negatively affect BMD, and while it may be too soon to state definitively, findings suggest that tea may even play a role in bone health.

- A study published recently in the *American Journal of Clinical Nutrition* found that older women who drank tea had higher BMD measurements than those who did not drink tea.⁸⁹
- Another recent study found that habitual tea-drinking was seen to have a significant beneficial effect on the BMD of adults (30 years and older), especially in those who had been habitual tea-drinkers for six or more years.⁹⁰
- Drinking tea is associated with preservation of hip structure in elderly women as assessed by dual-energy X-ray absorptiometry (DXA) areal bone mineral density (aBMD), an independent predictor for osteoporotic hip fracture, based on cross-sectional results (n=1027) and prospective analysis over 4 years (n=164) in a population of women, aged 70-85 years (p<0.05 for both results).⁹¹
- Studies in adolescent⁹² and postmenopausal women⁹³ found no relationship between caffeine intake and bone health.

###

03.15.09

References:

- ¹ Song WO, Chun OK. Tea is the major source of flavan-3-ol and flavonol in the U.S. diet. *J Nutr.* 2008;138:1543S–7S.
- ² Henning SM, Choo JJ, Heber D. Nongallated compared with gallated flavan-3-ols in green and black tea are more bioavailable. *J Nutr.* 2008;138:1529S–34S.
- ³ Auger C, Mullen W, Hara Y, Crozier A. Bioavailability of polyphenon E flavan-3-ols in humans with an ileostomy. *J Nutr.* 2008;138:1535S–42S.
- ⁴ Bolling BW, Chen CY, Blumberg JB. Tea and health: Preventive and therapeutic usefulness in the elderly? *Curr Opin Clin Nutr Metab Care.* 2009 Jan;12(1):42-8.
- ⁵ Weisburger JH. Tea and health: the underlying mechanisms. *Proc Soc Exp Biol Med* 1999;220:271-5.
- ⁶ Hertog MGL, Feskens EJM, Hollman PCH, et al. Dietary antioxidant flavonoids and risk of coronary disease: the Zutphen Elderly Study. *Lancet* 1993;342:1007-11.
- ⁷ Keli SO, Hertog MGL, Feskens EJM, Kromhout D. Dietary flavonoids, antioxidant vitamins, and incidence of stroke. *Arch Intern Med* 1996;156:637-42.
- ⁸ Duffy SJ, Keaney JF Jr, Holbrook M, Gokce N, Swerdloff PL, Frei B, Vita JA. Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation* 2001;104:151-6.
- ⁹ Isemura M, Saeki K, Kimura T, Hayakawa S, Minami T, Sazuka M. Tea catechins and related polyphenols as anti-cancer agents. *Biofactors.* 2000;13(1-4):81-5.
- ¹⁰ Mandel SA, Amit T, Kalfon L, Reznichenko L, Youdim MBH. Targeting multiple neurodegenerative diseases etiologies with multimodal-acting green tea catechins. *J Nutr.* 2008;138:1578S–83S.
- ¹¹ Kelly SP, Gomez-Ramirez M, Montesi JL, Foxe JJ. L-Theanine and caffeine in combination affect human cognition as evidenced by oscillatory alpha-band activity and attention task performance. *J Nutr.* 2008;138:1572S–7S.
- ¹² Sarkar, S., Sett, P., Chowdhury, T., and Ganguly, D.K. Effect of black tea on teeth. *J Indian Soc Pedod Prev Dent* 2000;18:139-140.
- ¹³ Hegarty VM, May HM, Khaw K-T. Tea drinking and bone mineral density in older women. *Am J Clin Nutr* 2000;71:1003-7.
- ¹⁴ Kamath AB, Wang L, Das H, Li L, Reinhold VN, Bukowski JF. Antigens in tea-beverage prime human Vgamma 2Vdelta 2 T cells in vitro and in vivo for memory and nonmemory antibacterial cytokine responses. *Proc Natl Acad Sci USA* 2003 May 13;100(10):6009-14.
- ¹⁵ Popkin BM, Armstrong LE, Bray GM, Caballero B, Frei B, Willett WC. A new proposed guidance system for beverage consumption in the United States. *Am J Clin Nutr.* 2006 Mar;83(3):529-42.
- ¹⁶ Hakim IA, Alsaif MA, Alduwaihy M, Al-Rubeaan K, Al-Nuaim AR, Al-Attas OS. Tea consumption and the prevalence of coronary heart disease in Saudi adults: results from a Saudi national study. *Prev Med* 2003;36(1):64-70.
- ¹⁷ Sesso HD, Gaziano JM, Buring JE, Hennekens CH. Coffee and tea intake and the risk of myocardial infarction. *Am J Epidemiol* 1999;149:162-7.
- ¹⁸ Mukamal KJ, Maclure M, Muller JE, Sherwood JB, Mittleman MA. Tea Consumption and Mortality After Acute Myocardial Infarction. *Circulation* 2002;105:2476.

-
- ¹⁹ Geleijnse JM, Launer LJ, Van der Kuip DA, Hofman A, Witteman JC. Inverse association of tea and flavonoid intakes with incident myocardial infarction: the Rotterdam Study. *Am J Clin Nutr*. 2002 May;75(5):880-6.
- ²⁰ Peters U, Poole C, Arab L. Does tea affect cardiovascular disease? A meta-analysis. *Am J Epidemiol* 2001;154(6):495-503.
- ²¹ Kuriyama S, Shimazu T, Ohmori K, Kikuchi N, Nakaya N, Nishino Y, Tsubono Y, Tsuji I. Green tea consumption and mortality due to cardiovascular disease, cancer, and all causes in Japan: the Ohsaki study. *JAMA*. 2006 Sep 13;296(10):1255-65.
- ²² Davies MJ, Judd JT, Baer DJ, Clevidence BA, Paul DR, Edwards AJ, Wiseman SA, Muesing RA, Chen SC. Black tea consumption reduces total and LDL cholesterol in mildly hypercholesterolemic adults. *J Nutr*. 2003 Oct;133(10):3298S-3302S.
- ²³ Vermeer MA, Mulder TP, Molhuizen HO. Theaflavins from black tea, especially theaflavin-3-gallate, reduce the incorporation of cholesterol into mixed micelles. *J Agric Food Chem*. 2008 Dec 24;56(24):12031-6.
- ²⁴ Geleijnse JM, Launer LJ, Hofman A, Pols HAP, Witteman JCM. Tea flavonoids may protect against atherosclerosis: the Rotterdam Study. *Arch Intern Med* 1999;159:2170-4.
- ²⁵ Duffy SJ, Keaney JF Jr, Holbrook M, Gokce N, Swerdloff PL, Frei B, Vita JA. Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation* 2001;104:151-6.
- ²⁶ Hodgson JM, Puddey IB, Burke V, Watts GF, Beilin LJ. Regular ingestion of black tea improves brachial artery vasodilator function. *Clin Sci* 2002;102(2):195-201.
- ²⁷ Hofmann CS, Sonenshein GE. Green tea polyphenol epigallocatechin-3 gallate induces apoptosis of proliferating vascular smooth muscle cells via activation of p53. *FASEB J*. 2003 Apr;17(6):702-4. Epub 2003 Feb 05.
- ²⁸ Ishikawa T, Suzukawa M, Ito T, Yoshida H, Ayaori M, Nishiwaki M, Yonemura A, Hara Y, Nakamura H. Effect of tea flavonoid supplementation on the susceptibility of low-density lipoprotein to oxidative modification. *Am J Clin Nutr* 1997;66:261-6.
- ²⁹ Vinson JA, Dabbagh YA, Serry MM, Jang J. Plant flavonoids, especially tea flavonols, are powerful antioxidants using an *in vitro* oxidation model for heart disease. *J Agric Food Chem* 1995;43:2800-2.
- ³⁰ Hirata K, Shimada K, Watanabe H, Otsuka R, Tokai K, Yoshiyama M, Homma S, Yoshikawa J. Black tea increases coronary flow velocity reserve in healthy male subjects. *Am J Cardiol*. 2004 Jun 1;93(11):1384-8, A6.
- ³¹ Vinson JA, Teufel K, Wu N. Green and black teas inhibit atherosclerosis by lipid, antioxidant, and fibrinolytic mechanisms. *J Agric Food Chem*. 2004 Jun 2;52(11):3661-5.
- ³² Yang YC, Lu FH, Wu JS, Wu CH, Chang CJ. The protective effect of habitual tea consumption on hypertension. *Arch Intern Med*. 2004 Jul 26;164(14):1534-40.
- ³³ Hakim IA, Chow HHS, Harris RB. Green tea consumption is associated with decreased DNA damage among GSTM1 positive smokers regardless of their hOGG1 genotype. *J Nutr*. 2008;138:1567S-71S.
- ³⁴ Hakim IA, Harris RB, Brown S, Chow HH, Wiseman S, Agarwal S, Talbot W. Effect of increased tea consumption on oxidative DNA damage among smokers: a randomized controlled study. *J Nutr*. 2003 Oct;133(10):3303S-3309S.
- ³⁵ Roy M, Chakrabarty S, Sinha D, Bhattacharya RK, Siddiqi M. Anticlastogenic, antigenotoxic and apoptotic activity of epigallocatechin gallate: a green tea polyphenol. *Mutat Res* 2003;523-524:33-41.
- ³⁶ Bhattacharyya A, Mandal D, Lahiry L, Sa G, Das T. Black tea protects immunocytes from tumor-induced apoptosis by changing Bcl-2/Bax ratio. *Cancer Lett*. 2004 Jun 25;209(2):147-54.

- ³⁷ Dora I, Arab L, Martinchik A, Sdvizhkov A, Urbanovich L, Weisgerber U. Black tea consumption and risk of rectal cancer in Moscow population. *Ann Epidemiol*. 2003 Jul;13(6):405-11.
- ³⁸ Su LJ, Arab L. Tea consumption and the reduced risk of colon cancer -- results from a national prospective cohort study. *Public Health Nutr*. 2002 Jun;5(3):419-25.
- ³⁹ Arts IC, Jacobs DR Jr, Gross M, Harnack LJ, Folsom AR. Dietary catechins and cancer incidence among postmenopausal women: the Iowa Women's Health Study (United States). *Cancer Causes Control*. 2002 May;13(4):373-82.
- ⁴⁰ Zheng W, Doyle TJ, Kushi LH, et al. Tea consumption and cancer incidence in a prospective cohort study of postmenopausal women. *Am J Epidemiol* 1996;144:175-81.
- ⁴¹ Sun CL, Yuan JM, Lee MJ, Yang CS, Gao YT, Ross RK, Yu MC. Urinary tea polyphenols in relation to gastric and esophageal cancers: a prospective study of men in Shanghai, China. *Carcinogenesis* 2002;23(9):1497-1503.
- ⁴² Ji BT, Chow WH, Hsing AW, McLaughlin JK, Dai Q, Gao YT, Blot WJ, Fraumeni JF Jr. Green tea consumption and the risk of pancreatic and colorectal cancers. *Int J Cancer*. 1997 Jan 27;70(3):255-8.
- ⁴³ Ohishi T, Kishimoto Y, Miura N, Shiota G, Kohri T, Hara Y, Hasegawa J, Isemura M. Synergistic effects of (-)-epigallocatechin gallate with sulindac against colon carcinogenesis of rats treated with azoxymethane. *Cancer Lett*. 2002 Mar 8;177(1):49-56.
- ⁴⁴ Lodovici M, Casalini C, De Filippo C, Copeland E, Xu X, Clifford M, Dolara P. Inhibition of 1,2-dimethylhydrazine-induced oxidative DNA damage in rat colon mucosa by black tea complex polyphenols. *Food Chem Toxicol*. 2000 Dec;38(12):1085-8.
- ⁴⁵ Isemura M, Saeki K, Kimura T, Hayakawa S, Minami T, Sazuka M. Tea catechins and related polyphenols as anti-cancer agents. *Biofactors*. 2000;13(1-4):81-5.
- ⁴⁶ Kuo PL, Lin CC. Green tea constituent (-)-epigallocatechin-3-gallate inhibits Hep G2 cell proliferation and induces apoptosis through p53-dependent and Fas-mediated pathways. *J Biomed Sci* 2003;10(2):219-27.
- ⁴⁷ Issa AY, Volate SR, et al. Green tea selectively targets initial stages of intestinal carcinogenesis in the AOM-ApcMin mouse model. [Carcinogenesis](#). 2007 Jul 17; [Epub ahead of print]
- ⁴⁸ Siddiqui IA, Saleem M. et al. Tea beverage in chemoprevention and chemotherapy of prostate cancer. [Acta Pharmacol Sin](#). 2007 Sep;28(9):1392-408.
- ⁴⁹ Hakim IA, Harris RB. Joint effects of citrus peel use and black tea intake on the risk of squamous cell carcinoma of the skin. *BMC Dermatol*. 2001;1(1):3. Epub 2001 Aug 01.
- ⁵⁰ Hakim IA, Harris RB, Weisgerber UM. Tea intake and squamous cell carcinoma of the skin: influence of type of tea beverages. *Cancer Epidemiol Biomarkers Prev*. 2000 Jul;9(7):727-31.
- ⁵¹ Lu YP, Lou YR, Lin Y, Shih WJ, Huang MT, Yang CS, Conney AH. Inhibitory effects of orally administered green tea, black tea, and caffeine on skin carcinogenesis in mice previously treated with ultraviolet B light (high-risk mice): relationship to decreased tissue fat. *Cancer Res* 2001 Jul 1;61(13):5002-9.
- ⁵² Ahmad N, Mukhtar H. Cutaneous photochemoprotection by green tea: a brief review. *Skin Pharmacol Appl Skin Physiol*. 2001 Mar-Apr;14(2):69-76.
- ⁵³ Katiyar SK, Bergamo BM, Vyalil PK, Elmets CA. Green tea polyphenols: DNA photodamage and photoimmunology. *J Photochem Photobiol B*. 2001 Dec 31;65(2-3):109-14.
- ⁵⁴ Katiyar SK, Perez A, Mukhtar H. Green tea polyphenol treatment to human skin prevents formation of ultraviolet light B-induced pyrimidine dimers in DNA. *Clin Cancer Res*. 2000 Oct;6(10):3864-9.

- ⁵⁵ Conney AH, Lu Y-P, Lou Y-R, Xie J-G, Huang M-T. Inhibitory effect of green and black tea on tumor growth. *Proc Soc Exp Biol Med* 1999;220:229-33.
- ⁵⁶ Li N, Zheng S, Han C, Chen J. The Chemoprotective Effects of Tea on Human Oral Precancerous Mucosa Lesions. *Proc Soc Exp Biol Med* 1999;220:218-24.
- ⁵⁷ Li N, Chen X, Liao J, Yang G, Wang S, Josephson Y, Han C, Chen J, Huang MT, Yang CS. Inhibition of 7,12-dimethylbenz[a]anthracene (DMBA)-induced oral carcinogenesis in hamsters by tea and curcumin. *Carcinogenesis* 2002;23(8):1307-13.
- ⁵⁸ Arts ICW. A review of the epidemiological evidence on tea, flavonoids, and lung cancer. *J Nutr*. 2008;138:1561S–6S.
- ⁵⁹ Yang G, Liu Z, Seril DN, et al. Black tea constituents, theaflavins, inhibit 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)-induced lung tumorigenesis in A/J mice. *Carcinogenesis* 1997;18:2361-5.
- ⁶⁰ Yang G, Wang Z-Y, Kim S, et al. Characterization of early pulmonary hyperproliferation and tumor progression and their inhibition by black tea in a 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis model with A/J mice. *Cancer Res* 1997;57:1889-94.
- ⁶¹ Ganguly C, Saha P, Panda CK, Das S. Inhibition of Growth, Induction of Apoptosis and Alteration of Gene Expression by Tea Polyphenols in the Highly Metastatic Human Lung Cancer Cell Line NCI-H460. *Asian Pac J Cancer Prev*. 2005 Jul-Sep;6(3):326-31.
- ⁶² Zhang M, Binns CW, Lee AH. Tea consumption and ovarian cancer risk: a case-control study in China. *Cancer Epidemiol Biomarkers Prev* 2002;11(8):713-8.
- ⁶³ Larsson SC, Wolk A. Tea consumption and ovarian cancer risk in a population-based cohort. *Arch Intern Med*. 2005 Dec 12-26;165(22):2683-6.
- ⁶⁴ Kumar N, Titus-Ernstoff L, Newcomb PA, Trentham-Dietz A, Anic G, Egan KM. Tea consumption and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev*. 2009 Jan;18(1):341-5.
- ⁶⁵ Kelly SP, Gomez-Ramirez M, Montesi JL, Foxe JJ. L-Theanine and caffeine in combination affect human cognition as evidenced by oscillatory alpha-band activity and attention task performance. *J Nutr*. 2008;138:1572S–7S.
- ⁶⁶ Egashira N, Ishigami N, et al. Theanine prevents memory impairment induced by repeated cerebral ischemia in rats. *Phytother Res*. 2007 Aug 17; [Epub ahead of print]
- ⁶⁷ Mandel SA, Amit T, Kalfon L, Reznichenko L, Youdim MBH. Targeting multiple neurodegenerative diseases etiologies with multimodal-acting green tea catechins. *J Nutr*. 2008;138:1578S–83S.
- ⁶⁸ Rezai-Zadeh K, Shytle D, Sun N, Mori T, Hou H, Jeanniton D, Ehrhart J, Townsend K, Zeng J, Morgan D, Hardy J, Town T, Tan J. Green tea epigallocatechin-3-gallate (EGCG) modulates amyloid precursor protein cleavage and reduces cerebral amyloidosis in Alzheimer transgenic mice. *J Neurosci*. 2005 Sep 21;25(38):8807-14.
- ⁶⁹ Hu G, Bidel S, et al. Coffee and tea consumption and the risk of Parkinson's disease. *Mov Disord*. 2007 Aug 21: [Epub ahead of print]
- ⁷⁰ Dulloo AG, Duret C, Rohrer D, Girardier L, Mensi N, Fathi M, Chantre P, Vandermander J. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *Am J Clin Nutr*. 1999 Dec;70(6):1040-5.
- ⁷¹ Chantre P, Lairon D. Recent findings of green tea extract AR25 (Exolise) and its activity for the treatment of obesity. *Phytotherapy* 2002;9(1):3-8.
- ⁷² Venables MC, Hulston CJ, Cox HR, and Jeukendrup AE. Green tea extract ingestion, fat oxidation, and glucose tolerance in healthy humans. *Am J Clin Nutr* 2008;87(3):778-84.

-
- ⁷³ Nagao T, Hase T and Tokimitsu I. A green tea extract high in catechins reduces body fat and cardiovascular risk in humans. *Obesity*. 2007 Jun;15:1473-83.
- ⁷⁴ Nagao T, Komine Y, Soga S, Meguro S, Hase T, Tanaka Y, Yokimitsu I. Ingestion of a tea rich in catechins leads to a reduction in body fat and malondialdehyde-modified LDL in men. *Am J Clin Nutr*. 2005 Jan;81(1):122-9.
- ⁷⁵ Murase T, Nagasawa A, Suzuki J, Hase T, Tokimitsu I. Beneficial effects of tea catechins on diet-induced obesity: stimulation of lipid catabolism in the liver. *Int J Obes Relat Metab Disord* 2002;26(11):1459-64.
- ⁷⁶ Murase T, Haramizu S, Shimotoyodome A, Tokimitsu I. Reduction of diet-induced obesity by a combination of tea-catechin intake and regular swimming. *Int J Obesity* 2005 Oct:1-8.
- ⁷⁷ Shimotoyodome A, Haramizu S, Inaba M, Murase T, Tokimitsu I. Exercise and green tea extract stimulate fat oxidation and prevent obesity in mice. *Med Sci Sports Exerc*. 2005 Nov;37(11):1884-92.
- ⁷⁸ Murase T, Haramizu S, Shimotoyodome A, Tokimitsu I, Hase T. Green tea extract improves running endurance in mice by stimulating lipid utilization during exercise. *Am J Physiol Regul Integr Comp Physiol*. 2006 Jun;290(6):R1550-6. renumber to 61
- ⁷⁹ Stote KS, Baer DJ. Tea consumption may improve biomarkers of insulin sensitivity and risk factors for diabetes. *J Nutr*. 2008;138:1584S-8S.
- ⁸⁰ Anderson RA, Polansky MM. Tea enhances insulin activity. *J Agric Food Chem* 2002;50(24):7182-6.
- ⁸¹ Kamath AB, Wang L, Das H, Li L, Reinhold VN, Bukowski JF. Antigens in tea-beverage prime human Vgamma 2Vdelta 2 T cells in vitro and in vivo for memory and nonmemory antibacterial cytokine responses. *Proc Natl Acad Sci U S A*. 2003 May 13;100(10):6009-14. Epub 2003 Apr 28.
- ⁸² Sarkar, S., Sett, P., Chowdhury, T., and Ganguly, D.K. Effect of black tea on teeth. *J Indian Soc Pedod Prev Dent* 2000;18:139-140.
- ⁸³ Yu, H., Oho, T., Xu, L. X. Effects of several tea components on acid resistance of human tooth enamel. *J Dent* 1995;13:101-105.
- ⁸⁴ Linke HA, LeGeros RZ. Black tea extract and dental caries formation in hamsters. *Int J Food Sci Nutr* 2003;54(1):89-95.
- ⁸⁵ Bassiouny MA, Kuroda S, Yang J. 2008. Topographic and radiographic profile assessment of dental erosion. Part III: Effect of green and black tea on human dentition. *General Dentistry*. Jul-Aug;56(5):451-61.
- ⁸⁶ Kushiyama M, Shimazaki Y, Murakami M, Yamashita Y. 2009. Relationship between intake of green tea and periodontal disease. *J Periodontol*. Mar;80(3):372-7.
- ⁸⁷ Curhan GC, Willett WC, Speizer FE, Stampfer MJ. Beverage use and risk of kidney stones in women. *Ann Intern Med* 1998;128:534-40.
- ⁸⁸ Curhan GC, Willett WC, Rimm EB, Spiegelman D, Stampfer MJ. Prospective study of beverage use and the risk of kidney stones. *Am J Epidemiol* 1996;143:240-7.
- ⁸⁹ Hegarty VM, May HM, Khaw K-T. Tea drinking and bone mineral density in older women. *Am J Clin Nutr* 2000;71:1003-7.
- ⁹⁰ Wu CH, Yang YC, Yao WJ, Lu FH, Wu JS, Chang CJ. Epidemiological evidence of increased bone mineral density in habitual tea drinkers. *Arch Intern Med*. 2002 May 13;162(9):1001-6.
- ⁹¹ Devine A, Hodgson JM, Dick IM, Prince RL. Tea drinking is associated with benefits on bone density in older women. *Am J Clin Nutr*. 2007;86(4):1243-7.

⁹² Lloyd T, Rollings NJ, Kieselhorst K, Egli DF, Mauger E. Dietary caffeine intake is not correlated with adolescent bone gain. *J Am Coll Nutr* 1998;17:454-7.

⁹³ Lloyd T, Johnson-Rollings N, Egli DF, Kieselhorst K, Mauger EA, Cusatis DC. Bone status among postmenopausal women with different habitual caffeine intakes: a longitudinal investigation. *J Am Coll Nutr* 2000;19:256-61